

Meet the suPARnostic® POC+

The point-of-care solution for measuring suPAR

suPARnostic®
by ViroGates

- Testing possible without laboratory involvement
- Ease of use
- Simple finger prick test
- Results available in 20 minutes

The suPARnostic® POC+ product is a point-of-care finger prick test which detects the level of suPAR in the blood.

The suPARnostic® POC+ has full CE-IVD accreditation for use in professional healthcare settings to identify inflammation and immune activation.

The suPARnostic® POC+ is based on the GENSPEED technology that allows quick and fully automated processing of finger prick blood samples. The test takes approximately 20 minutes.

The **suPARnostic® POC+** is useful in acute care and in the pre-hospital settings.

A kit contains ready-to-use reagents sufficient to perform 25 tests.

Contact our experts about your needs and they will help you.

Contact us as sales@virogates.com



CSO at ViroGates Jesper Eugen-Olsen, PhD, says: "With the POC+, we can finally measure the suPAR level in a drop of blood from the finger. This allows for easy determination of chronic inflammation, disease severity, and prognosis in settings without laboratories."

Read more about the
suPARnostic® POC+



suPARnostic® POC+ is quick and easy to



use

The size and technology used in suPARnostic® POC+ ensure excellent accessibility and mobility of the device. The small footprint means that it fits into smaller spaces and can be carried between multiple departments.

Test benefits

- Empower the clinical staff
- 22% reduction in hospital admissions
- Reduce healthcare costs



1

Add a finger prick blood drop to the test cassette



2

Place the sample in the POC+ device



3

Wait 17 minutes



4

The result is available on the screen



5

Clinical decision making is supported

